

Front

70.00 mm	
Patient Alert Card Apixaban 2.5 mg Film-coated Tablets Apixaban 5 mg Film-coated Tablets This medicine will be referred to as Apixaban tablets throughout this Patient Alert Card. Carry this card with you at all times Show this card to your pharmacist, dentist and any other healthcare professionals that treat you. Information for patients • Take Apixaban tablets regularly as instructed. If you miss a dose, take it soon as you remember and continue to follow your dosing schedule. • Do not stop taking Apixaban tablets	without talking to your doctor, as you are at risk of suffering from a stroke or other complications. • Apixaban tablets help to thin your blood. However, this may increase your risk of bleeding. • Signs and symptoms of bleeding include bruising or bleeding under the skin, tar-colored stools, blood or urine, nosebleed, dizziness, tiredness, paleness or weakness, sudden severe headache, coughing up blood or vomiting blood. • If the bleeding does not stop on its own, seek medical attention immediately. • If you need surgery or any invasive procedure, inform your doctor that you are taking Apixaban tablets.
40.00 mm	

Back

I am under anticoagulation treatment with Apixaban tablets to prevent blood clots Information for healthcare professionals • Apixaban tablets are an oral anticoagulant acting by direct selective inhibition of factor Xa. • Apixaban tablets may increase the risk of bleeding. In case of major bleeding events, it should be stopped immediately. • Treatment with Apixaban tablets does not require routine monitoring of exposure. A calibrated quantitative anti-Factor Xa assay may be useful in exceptional situations, e.g. overdose and emergency surgery (prothrombin time (PT), international	normalized ratio (INR) and activated partial thromboplastin time (aPTT) clotting tests are not recommended) – see SmPC. • An agent to reverse the anti-Factor Xa activity of Apixaban tablets is available. Please complete this section or ask your doctor to do it Name: _____ Birth Date: _____ Indication: _____ Dose: _____ mg twice daily Doctor's telephone: _____
40.00 mm	
70.00 mm 	

Minimum font size: 5 pt